

DOCUMENTATION, ANTI-AGING ACTIVITIES AND PHYTOCHEMICAL
PROFILING OF SELECTED MEDICINAL PLANTS USED BY JAKUN
WOMEN IN KAMPUNG PETA, MERSING, JOHOR

NUR AMALINA BINTI ISMAIL

A thesis submitted in
fulfillment of the requirement for the award of the degree of
Master of Science

Faculty of Applied Science and Technology
Universiti Tun Hussein Onn Malaysia

SEPTEMBER 2017

DEDICATION

For my *abah* (En. Ismail Bin Ebrahim) and *mama* (Pn Siti Habsah Bt. Hashim),
who always inspired me with their hardwork and successs,
who always love, pray and give endless support to their children.

And also for my lovely siblings (Izzati, Nabila & Amir) for their constant motivation
and support along this journey.

May Allah reward them with Jannah



PTTA UTHM
PERPUSTAKAAN TUNKU TUN AMINAH

ACKNOWLEDGEMENT

Alhamdulillah. All praise be to Allah the Almighty for his blessings. I wish to express my sincere appreciation to my main supervisor, Assoc. Prof. Dr. Mohd. Fadzelly Bin Abu Bakar, for his guidance, support and encouragement since the undergraduate years till the postgraduate journey. I also wish to thank my co-supervisor, Prof. Datin Dr. Maryati Mohamed and Dr. Faridah Kormin for their guidance and precious knowledge on traditional knowledge and experimental design. I truly appreciate their endeavour in instilling the good character in me to become better researcher.

I also would like to thank Assoc. Prof. Dr. Abdah Bt. Akim for supporting me during my research attachment in Biomedical Laboratory, Faculty of Medicine and Health Science, Universiti Putra Malaysia (UPM), Mr. Azizul Isha from Laboratory of Natural Products, UPM, Department of Orang Asli Development (JAKOA) and Johor National Parks Corporation (PTNJ) for the approval to conduct the research and not to forget, Mr. Ishak Ayob and Mdm. Siti Nazmilah Misrin for technical assistance in Food Microbiology Lab.

I would like to give special appreciation for my research colleague from Centre of Research for Sustainable Uses of Natural Resources (CoR-SUNR), for their endless support throughout this journey. My deepest gratitude to Jakun community of Kampung PETA especially the key informants who had given their cooperation and shared their precious knowledge during interview session. Last but not least, this thesis would not be complete without the financial support from the Public Service Department (JPA) under *Program Mahasiswa Cemerlang* (PMC) scholarship, Fundamental Research Grant Scheme (FRGS) No. 1435, FRGS No. 1560, and Knowledge Transfer Programme 1568 (KTP 1568) Grant Scheme.

ABSTRACT

Traditional knowledge of indigenous people could become the baseline information for the discovery of anti-aging agent. The objectives of this study were to document the knowledge of Jakun people in Kampung Peta, Mersing, Johor on medicinal plants for women's healthcare; to investigate the optimal formulations of herbal mixture used by Jakun women based on phytochemicals content and antioxidant activity; to determine the anti-aging potential of the selected formulations; and to investigate the major phytochemical constituents in the formulations. Based on qualitative analysis from semi-structured interview, twelve species of medicinal plants have been documented for women's healthcare. Among species documented, four species, *Cnestis palala* (Pengesep), *Urceola micrantha* (Serapat), *Labisia pumila* (Kacip fatimah) and *Microporus xanthopus* (Kulat kelentit kering) that were prepared in the form of mixture have been used for formulation study. About 24 formulations have been developed from the simplex centroid design and tested for total phenolic content (Folin-Ciocalteu method), total flavonoid content (aluminium chloride colorimetric method) and three different antioxidant assays (DPPH scavenging, ABTS decolourization and FRAP assays). Single formulation of *Cnestis palala*, single formulation of *Urceola micrantha* and binary mixture of *C. palala* and *U. micrantha* are among the optimal formulations with high phytochemicals content and antioxidant activities that were further evaluated for anti-aging activities. For anti-aging activities, five enzymatic assays have been tested on the three formulations which are matrix metalloproteinase-1 (MMP-1) inhibition, elastase inhibition, tyrosinase inhibition, acetyl- and butyrylcholinesterase inhibition assays. Single formulation of *U. micrantha* showed the highest inhibition towards MMP-1 (49.44 ± 4.11 %) and elastase enzymes (20.33 ± 2.52 %), while single formulation of *C. palala* showed highest inhibitions towards tyrosinase (14.06 ± 0.31 %), acetylcholinesterase (32.92 ± 2.13 %) and butyrylcholinesterase (34.89 ± 2.84 %) enzymes. The identification of phytochemicals compound have been carried out using gas chromatography-mass spectrometer (GC-MS), which showed the presence of 2,2-dimethoxybutane and 2,3-dihydro-3,5-dihydroxy-6-methyl-4H-pyran-4-one (DDMP) in the three formulations extract. The presence of catechol and quinic acid in *U. micrantha* extract might possibly contribute to anti-aging activities of the extract. These findings could become baseline for the exploration of novel anti-aging agents from natural source by using the traditional knowledge of indigenous people.

ABSTRAK

Pengetahuan tradisi orang asli boleh menjadi maklumat asas kepada penemuan agen anti-penuaan. Objektif kajian ini adalah untuk mendokumenkan pengetahuan orang Jakun di Kampung Peta, Mersing, Johor terhadap tumbuhan ubatan yang digunakan untuk penjagaan kesihatan wanita; untuk mengkaji formulasi optimum bagi campuran herba yang digunakan oleh wanita Jakun berdasarkan kandungan fitokimia dan aktiviti antioksidasi; untuk menentukan potensi anti-penuaan bagi formulasi terpilih; dan untuk mengkaji komposisi fitokimia dalam formulasi. Berdasarkan analisis kualitatif dari temu bual separa berstruktur, dua belas spesies tumbuhan ubatan telah didokumenkan bagi penjagaan kesihatan wanita. Dari spesies yang telah didokumenkan, empat spesies, *Cnestis palala* (Pengesep), *Urceola micrantha* (Serapat), *Labisia pumila* (Kacip Fatimah) dan *Microporus xanthopus* (Kulat kelentit kering) yang di sediakan dalam bentuk campuran digunakan bagi kajian formulasi. Sekitar 24 formulasi telah dibangunkan menggunakan reka bentuk simpleks sentroid dan diuji untuk jumlah kandungan fenolik (kaedah Folin-Ciocalteu), jumlah kandungan flavonoid (kaedah kolorimetrik aluminium klorida) dan tiga asai antioksidasi (asai penghapusan DPPH, penyahwarnaan ABTS dan FRAP). Formulasi tunggal *Cnestis palala*, formulasi tunggal *Urceola micrantha* dan campuran perduaan *C. palala* dan *U. micrantha* adalah antara formulasi optimal dengan kandungan fitokimia dan aktiviti antioksidasi yang tinggi, yang dinilai bagi aktiviti anti-penuaan. Bagi aktiviti anti-penuaan, lima asai enzim telah diuji bagi tiga formulasi, iaitu asai perencatan matriks metalloproteinase-1 (MMP-1), perencatan elastase, perencatan tirosinase, perencatan asetil- dan butirilkolinesterase. Formulasi tunggal *U. micrantha* menunjukkan perencatan yang tinggi terhadap MMP-1 ($49.44 \pm 4.11\%$) dan enzim elastase ($20.33 \pm 2.52\%$ perencatan), manakala formulasi tunggal *C. palala* menunjukkan perencatan yang tinggi terhadap enzim-enzim tirosinase ($14.06 \pm 0.31\%$), asetilkolinesterase ($32.92 \pm 2.13\%$) dan butirilkolinesterase ($34.89 \pm 2.84\%$). Pengenalpastian sebatian fitokimia telah dijalankan menggunakan kromatografi gas-spektrometri jisim menunjukkan kehadiran 2,2-dimetoksibutane and 2,3-dihidro-3,5-dihidroksi-6-metil-4H-pyran-4-one (DDMD) di dalam tiga ekstrak formulasi. Kehadiran katekol dan asid kuinik dalam ekstrak *U. micrantha* berkemungkinan dapat menyumbang aktiviti anti-penuaan bagi ekstrak. Penemuan ini dapat menjadi asas kepada penerokaan agen anti-penuaan yang baharu dari sumber semulajadi dengan menggunakan pengetahuan tradisi orang asli.

TABLE OF CONTENTS

TITLE	i
DECLARATION	ii
DEDICATION	iii
ACKNOWLEDGEMENT	iv
ABSTRACT	v
ABSTRAK	vi
TABLE OF CONTENTS	vii
LIST OF TABLES	xii
LIST OF FIGURES	xiii
LIST OF SYMBOLS AND ABBREVIATIONS	xiv
LIST OF APPENDICES	xvi
LIST OF PUBLICATIONS	xvii
CHAPTER 1 INTRODUCTION	1
1.1 Background of study	1
1.2 Problem statement	5
1.3 Objectives	5
1.4 Significance of study	6
CHAPTER 2 LITERATURE REVIEW	7
2.1 Documentation of traditional knowledge	7
2.1.1 Definition of terms	7
2.1.2 The importance of documenting the traditional knowledge	8
2.1.2.1 Conserving genetic resources and natural heritage	8
2.1.2.2 Intellectual property and protection against biopiracy	9
2.1.2.3 Foundation to the drug discovery	10

2.1.3	Issues in traditional knowledge and ethnobotany	10
2.2	Jakun community	11
2.3	Ethnomedicinal knowledge of plants	12
2.3.1	Ethnomedicinal knowledge of plants by the Jakun community	12
2.3.2	Medicinal plants for women's healthcare and beauty	12
2.4	Aging	13
2.4.1	Terminologies	13
2.4.2	Theories of aging	14
2.4.3	Role of oxidation in aging	15
2.4.4	Aging through skin	15
2.4.4.1	Mechanism of wrinkle formation	15
2.4.4.2	Mechanism of hyperpigmentation	17
2.4.5	Alzheimer's disease and its mechanism	17
2.5	Intervention of aging	18
2.5.1	Antioxidant and its mechanism	18
2.5.2	Anti-wrinkle	20
2.5.3	Anti-hyperpigmentation	20
2.5.4	Cholinesterase inhibition	21
2.6	Medicinal plants and anti-aging activities	21
2.6.1	Medicinal plants with antioxidant potential	22
2.6.2	Medicinal plants with anti-skin aging properties	23
2.6.2.1	Medicinal plants as anti-wrinkle agent	23
2.6.2.2	Medicinal plants as anti-tyrosinase agent	24
2.6.3	Medicinal plants as anti-Alzheimer agent	25
2.7	Phytochemicals and anti-aging activities	26
2.7.1	Phenolics	26

2.7.2	Flavonoids	27
2.7.3	Stilbenoids	28
2.7.4	Carotenoids	29
2.7.5	Terpenoids	29
2.8	Herbal mixture	29
2.8.1	Application of mixture design in medicinal plants formulation	30
2.8.2	Synergism of herbal mixture	30
CHAPTER 3	METHODOLOGY	32
3.1	Chemicals and reagents	32
3.2	Equipments and apparatus	33
3.3	Study design and sampling site	33
3.4	Selection of key informants	35
3.5	Ethnobotanical study of women's medicinal plants	36
3.5.1	Ethical authorization	36
3.5.2	Semi-structured interview	36
3.5.3	Data analysis	37
3.6	Plant materials	
3.6.1	Identification of plant species	37
3.6.2	Criteria for selection of medicinal plants.	37
3.6.3	Collection and preparation of raw Material	38
3.7	Mixture design	39
3.7.1	Simplex centroid mixture design	39
3.7.2	Validation of optimal formulations	40
3.8	Preparation of sample infusion	42
3.9	Phytochemical analysis	42
3.9.1	Total phenolic content	42
3.9.2	Total flavonoid content	42
3.10	Antioxidant activities	43

3.10.1	2, 2'-azino-bis(3-ethylbenzothiazoline-6-sulphonic acid (ABTS) cation radical decolorization assay	43
3.10.2	2, 2-diphenyl-1-picryl-hydrazyl (DPPH) free radical scavenging activity	43
3.10.3	Ferric reducing/antioxidant power (FRAP) assay	44
3.11	Anti-wrinkle assay	44
3.11.1	Collagenase inhibition assay	44
3.11.2	Elastase inhibition assay	45
3.12	Anti-hyperpigmentation assay	45
3.12.1	Tyrosinase inhibition assay	45
3.13	Anti-Alzheimer	46
3.13.1	Acetylcholinesterase inhibition assay	46
3.13.2	Butyrylcholinesterase inhibition assay	46
3.14	Profiling of chemical compound by using Gas Chromatography Mass Spectrometry	47
3.15	Statistical analysis	47
CHAPTER 4	RESULT AND DISCUSSION	48
4.1	Ethnobotanical knowledge of medicinal plants for women's health	48
4.1.1	Plant species, families, and growth habits	48
4.1.2	Uses of plants for the women's healthcare	51
4.1.3	Parts used	51
4.1.4	Preparation and administration	52
4.1.5	Frequency of citation	54
4.1.6	Selection of medicinal plants for bioassay	54
4.2	Applications of simplex centroid mixture design for optimization of herbal mixture	55
4.2.1	Evaluation of phytochemicals content	57
4.2.1.1	Total phenolic content	57
4.2.1.2	Total flavonoid content	59



4.2.2	Evaluation of antioxidant activities	60
4.2.2.1	ABTS scavenging activity	61
4.2.2.2	DPPH radical scavenging activity	62
4.2.2.3	FRAP reducing ability	63
4.2.3	Effect of single, binary, ternary and quaternary polyherbal formulations on the phytochemicals content and antioxidant activities	64
4.2.4	ANOVA analysis for phytochemicals and antioxidant activities	65
4.2.5	Synergistic and antagonistic effects of herbal mixture on phytochemicals and antioxidant activities	67
4.2.6	Surface plots	71
4.2.7	Optimization of polyherbal formulations	73
4.3	Anti-aging activities of optimal formulations	75
4.3.1	Anti-collagenase activity of selected formulations	76
4.3.2	Anti-elastase activity	78
4.3.3	Anti-tyrosinase activity	80
4.3.4	Anti-cholinesterase activity	82
4.4	Correlation analysis between phytochemicals content and anti-aging activities	86
4.5	Phytochemicals profiling of selected formulations	90
CHAPTER 5	CONCLUSION	95
5.1	General conclusion	95
5.2	Limitation of the study	96
5.3	Recommendation for future research	96
	REFERENCES	98
	APPENDICES	122

LIST OF TABLES

2.1	Ethnomedicinal plants used by indigenous people in Peninsular Malaysia	13
3.1	Characteristics of selected key informants in Kampung Peta	36
3.2	List of plants selected for further analysis	37
3.3	Experimental design layout for herbal formulations	41
4.1	Ethnobotanical information of medicinal plants used for women's healthcare	49
4.2	Experimental design layout and the responses for herbal formulations	56
4.3	Average of TPC, TFC, ABTS, DPPH, and FRAP in different polyherbal formulations	65
4.4	ANOVA table of phytochemicals (TPC and TFC) and antioxidant activities (ABTS, DPPH and FRAP)	66
4.5	Coefficients of each model fitted and their level of significance determined by p-value.	70
4.6	Formulation selection based on desirability function	74
4.7	Predicted and experimental value of optimal formulations	75
4.8	Elastase inhibition activity of samples and elastatinal (standard)	79
4.9	Correlation between the phytochemicals content, antioxidant activities with anti-aging activities.	87
4.10	Major chemical constituents identified in the water extract of <i>C. palala</i> by GC-MS	91
4.11	Major chemical constituents identified in the water extract of <i>U. micrantha</i> by GC-MS	92
4.12	Chemical composition of binary combination of <i>C. palala</i> and <i>U. micrantha</i>	92

LIST OF FIGURES

3.1	Flowchart of the study design	34
3.2	Location of Kampung Peta and Taman Negara Johor Endau Rompin	35
3.3	Picture displaying selected medicinal plants. (a) <i>Cnestis palala</i> (b) <i>Urceola micrantha</i> (c) <i>Labisia pumila</i> (d) <i>Microporus xanthopus</i>	38
3.4	Arrangement of the simplex centroid design with four	39
4.1	Percentage of species based on plant parts used	52
4.2	Percentage of species for each method of preparation	53
4.3	Percentage of species based on the way of administration	54
4.4	Surface plots (a) total phenolic content [TPC], (b) total flavonoid content [TFC], (c) 2, 2'-azino-bis (3- ethylbenzothiazoline-6-sulphonic acid) scavenging activity [ABTS], (d) 2, 2-diphenyl-1-picryl-hydrazyl free radical scavenging activity [DPPH], (e) ferric reducing/antioxidant power activity [FRAP]	72
4.5	MMP-1 inhibition of samples and standard	77
4.6	Tyrosinase inhibition of optimal formulations	81
4.7	Acetylcholinesterase inhibition activity of tested formulations	84
4.8	Butyrylcholinesterase inhibition activity of samples and standard	85
4.9	Correlation between phytochemicals and antioxidant activities of 24 formulations	88
4.10	Gas Chromatograms of (a) hot water extract of <i>C. palala</i> , (b) hot water extract of <i>U. micrantha</i> , and (c) hot water extract of binary combination of <i>C. palala</i> and <i>U. micrantha</i>	90
4.11	Similar compounds found in all formulation extracts	93

LIST OF SYMBOLS AND ABBREVIATION

°C	-	Celsius
h	-	Hour
L	-	Liter
mL	-	Milliliter
mg	-	Milligram
min	-	Minute
μL	-	Microliter
μM	-	Micromolar
μU	-	Microunit
nm	-	Nanometer
R ²	-	Coefficient of determination
Aβ	-	Amyloid Beta
ABS	-	Access and benefit sharing
ABTS	-	2,2'-Azino-bis(3-ethylbenzothiazoline-6-sulphonic acid
AChE	-	Acetylcholinesterase enzyme
AD	-	Alzheimer's Disease
AP-1	-	Activator protein-1
BuChE	-	Butyrylcholinesterase enzyme
CaCl ₂	-	Calcium chloride
CBD	-	Convention on Biological Diversity
CE	-	Catechin equivalent
DMSO	-	Dimethylsulfoxide
DNA	-	Deoxyribonucleic acid
DOE	-	Design of Experiment
DDMP	-	2,3-dihydro-3,5-dihydroxy-6-methyl-4H-pyran-4-one
DPPH	-	2,2-diphenyl-1-picrylhydrazyl
DTNB	-	5,5'-dithio-bis-[2-nitrobenzoic acid]
ECM	-	Extracellular matrix

FDA	-	Food and Drug Administration
FRAP	-	Ferric-reducing antioxidant power
GAE	-	Gallic acid equivalent
GC-MS	-	Gas Chromatography- Mass Spectrometry
HEPES	-	4-(2-hydroxyethyl)-1-piperazineethanesulfonic acid
IK	-	Indigenous knowledge
JAKOA	-	Department of Orang Asli Development
L-DOPA	-	L-dihydroxyphenylalanine
MAPK	-	Mitogen-activated protein kinase
MMP	-	Matrix metalloproteinase
MRNA	-	Messenger ribonucleic acid
NNGH	-	N-Isobutyl-N-(4-methoxyphenylsulfonyl) glycyldihydroxamic
PIC	-	Prior Informed Consent
PTNJ	-	Johor National Park Corporation
ROS	-	Reactive oxygen species
TBARS	-	Thiobarbituric acid reactive substance
TEAC	-	Trolox equivalent antioxidant capacity
TFC	-	Total flavonoid content
TK	-	Traditional Knowledge
TMK	-	Traditional Medicinal Knowledge
TNJER	-	Taman Negara Johor Endau Rompin
TPC	-	Total phenolic content
TPTZ	-	2,4,6-tripyridyl-S-triazine
TRP-1	-	Tyrosinase related protein 1
UV	-	Ultra violet
VMS	-	Volatile methylsiloxane
WHO	-	World Health Organization

LIST OF APPENDICES

APPENDIX	TITLE	PAGE
A	Research approval from PTNJ	122
B	Research approval from JAKOA	123
C	Prior Informed Consent	124
D	Semi structured questionnaire	126
E	Gallic acid standard curve	128
	Catechin standard curve	
F	Ascorbic acid standard curve	129
	Ferrous sulphate standard curve	
G	Normal plot of residual for TPC and TFC	130
H	Normal plot of residual for antioxidant activities	131

LIST OF PUBLICATIONS

Journals :

- i. **Ismail, N. A.**, Abu Bakar, M. F., Kormin, F., Linatoc, A. C., Maryati, M. 2017. Application of statistically mixture simplex-centroid design to optimize the TPC and TFC on the proportion of polyherbal formulation used by Jakun women. *Journal of Engineering & Applied Science* (SCOPUS). [Accepted].
- ii. **Ismail, N. A.**, Abu Bakar, M. F., Kormin, F., Maryati, M., Akim, A., Isha, A. Anti-aging potential and phytochemical screening of optimal formulations from selected medicinal plants used by Jakun women in Kg. Peta, Johor. *Oxidative Medicine and Cellular Longevity* (SCOPUS). [Submitted]

Proceedings/ Conference/Seminars :

- i. **Ismail, N. A.**, Maryati, M. & Abu Bakar, M. F. (2015). Medicinal Plants Used For Women's Healthcare Among The Jakun Community In Kg. Peta : A Preliminary Study. In O. Asiah, H. Lim, B. Chee, M. N. Musaadah, M. Mastura, K. Getha, S. Vimala (Eds.), *Proceeding FRIM No. 7 Persidangan Industri Herba* (pp. 303–308). Putrajaya: Forest Research Institute Malaysia

CHAPTER 1

INTRODUCTION

1.1 Background of study

Aging is inevitable and considered as natural biological process. However, the changes in physiology of human due to aging are undesirable. Aging is physiologically characterized as a continuous, generalized systemic organ dysfunction, which lead to the increased vulnerability to environmental challenge and increase the tendency to suffer diseases and death. Generally, the incidence of degenerative diseases such as osteoporosis, cardiovascular disease, type 2 diabetes, cancer and Alzheimer's disease escalated with increasing age (Si & Liu, 2014) .

Physically, the obvious sign of aging could be seen on the skin with the formation of wrinkles and atypical pigmentation (Masaki, 2010). The skin condition such as epidermal thinning, reduction in dermal collagen content, decreased in laxity diminished skin moisture, and impaired wound healing have been reported in post-menopausal women (Irrera *et al.*, 2017). Exposure of skin to extrinsic factors (environmental aggressor such as UV radiation and smoking) is in the 4th rank that contribute to the nonfatal disease burden and the effects arise as humans get older (Tobin, 2017).

Brain also would be affected as person aged. Normal aging will cause decline in brain function and increase the tendency to suffer from neuronal degeneration. The decline in brain cholinergic system, that contribute to memory function will eventually lead to various age-related neurodegenerative diseases such as Parkinson's disease, Alzheimer's disease (AD), multiple sclerosis, Huntington disease and amyotrophic lateral sclerosis (Solanki *et al.*, 2016). Other than individual genetic factors, the external factors such as nutrition, smoking, alcohol,

environmental conditions could contribute to the aging in human. Oxidative stress has been identified to be one of the major factors that accelerates the aging process (Rahal *et al.*, 2014). Oxidation will cause oxidative injury to the neuron and skin structure. Antioxidant compounds are able to defend human body from oxidation process. Eventhough human body contains its own self defense system, it is insufficient to fight against the cumulative damage due to reactive oxygen species (ROS).

Plants are known to contain numerous bioactive compounds which could potentially act as antioxidant with anti-wrinkle and anti-Alzheimer's properties that protect the skin against key enzyme in wrinkle formation and brain from aging (Nema *et al.*, 2011). The exploration of medicinal plants as potential for anti-aging supplement or drugs are rapidly growing for the past few years. According to Global Industry Analyst, nutricosmetic market is estimated to reach US\$ 7.4 billion by 2020, driven by "beauty from within" trend (Analyst, 2015). This indicates that people worldwide are constantly searching for the intervention with nutraceutical benefit that could lead to the improvement and maintenance of their appearances and internal health. The nutricosmetic is the result of convergence between nutraceutical and cosmaceutical field with major claim as an anti-aging (Anunciato & Filho, 2012). The examples of nutricosmetic products include "beauty pills", tablets, liquid (i.e herbal tea), granulates or foods formulation which is believe to be able to reduce wrinkle formation by fighting free radicals generated by solar radiation (Taeymans *et al.*, 2014).

More than 89 medicinal plants have been used for cosmetic applications in Malaysia, such as *Centella asiatica* (Apiaceae), *Cosmos caudatus* (Asteraceae), *Curcuma xanthorrhiza* (Zingiberaceae) and *Ficus deltoidea* (Moraceae) are traditionally known to preserve youthful appearance of women (Narayanaswamy & Ismail, 2015). Usually, these plants contain antioxidant with anti-wrinkle properties. In spite of the wrinkle inhibition, the antioxidant also possess cholinesterase inhibitory activities which is one of the treatment in AD prevention. In a review by Natarajan *et al.* (2013) on Asian plants, more than 40 herbal remedies have been used traditionally and being proven scientifically to possess anti-inflammatory, anti-cholinesterase and antioxidant activities, which are contributed by its bioactive compounds such as alkaloids, flavonoids, steroids, saponins, terpenoids and essential oils. Despite the progress made in aging and Alzheimer's disease research in the last

decades, no treatment with a strong disease-modifying effect is currently available (Natarajan *et al.*, 2013).

Traditional knowledge (TK) on medicinal plants has been one of the sources in drug discovery from natural sources. TK has been defined as the knowledge, innovations and practices of indigenous and local communities around the world which were developed from experience gained over the centuries and adapted to local culture and environment, and transmitted orally from generation to generation (MoNRE, 2012). Traditional Medicinal Knowledge (TMK) is one of the sub categories of TK, other than traditional agricultural and ecological knowledge (Van Overwalle, 2005). According to World Health Organization (WHO, 2003), about 80% of people around the world utilize traditional medicine as their primary healthcare whereas 65% of citizen in developing nation use traditional medicine as an alternative to their healthcare maintenance. In National Policy on Biological Diversity 2016-2025, TK has been highlighted to be one of the key elements in conservation and sustainable uses of biodiversity (MoNRE, 2012). In addition, Malaysia has implemented an act on Access to Biological Resources and Benefit sharing bill 2017 in order to preserve the traditional knowledge of local and indigenous people, and also to ensure the fair and equitable sharing from the knowledge provided by the indigenous people.

Malaysia is ranked 12th among countries of the world with megabiodiversity and at least 95 subgroups of indigenous people that make up Malaysia's multiracial population (Masron *et al.*, 2013). The wonderful knowledge and respect towards plants have originated from indigenous people, usually called 'Orang Asli', who inherited the knowledge and practices from their ancestors (Adnan & Othman, 2012). The indigenous and local communities usually live in an area where majority of plants genetic resources are found. In this regard, the preservation of the TK through documentation is crucial to maintain the biodiversity in Malaysia (MonRE, 2012). However, TK suffer from erosion and biopiracy. These documentation is the utmost defensive approach for conserving the TK from erosion, acquisition and exploitation by third parties (Van Overwalle, 2005).

Jakun, Orang Kuala, Orang Kanaq, Orang Seletar and Temuan are ethnic groups recognized as orang asli that live in the southern part of Malaysia, Johor (Masron *et al.*, 2013). The Jakun tribe, who live in Kampung Peta, Endau, Mersing, Johor are the holder and users of TK. Despite modernization of this country, most of

the older generations in Jakun community still rely on traditional medicine for their healthcare. They utilize plants and animals to treat various ailments. The earliest record of ethnomedicinal knowledge of Jakun on plants has been documented by Taylor & Wong (1987), whereas the recent documentation of plant by Jin (2005) include the use of palm for medicinal purposes, Ismail *et al.* (2015) on the medicinal plants used for malarial treatment and Sabran *et al.* (2016) on medicinal plants used for treatment of tuberculosis. However, no study has been focusing on the use of medicinal plants specifically for maintenance of women's healthcare in Jakun community. The Convention on Biological Diversity (CBD) has recognized the role of women and local communities in conservation of biological diversity. In fact, a review on medicinal plants used for women healthcare in South-East Asia revealed 2000 different plant species have been used in 5000 combinations to treat various ailments for women (Boer & Cotingting, 2014). There is a clear need to document the TK of women in Jakun community on the use of medicinal plants, particularly in health and beauty maintenance.

While taking into account the individual medicinal plants for the treatment of disease, the consumption of traditional herbs in the form of mixture is believed to exert higher pharmacological and therapeutic efficacy compared to the single medicinal plants (Guimarães *et al.*, 2011; Wang *et al.*, 2014). Few reported studies showed that mixture of medicinal plants exhibit higher bioactivities compared to the single plants (Xu *et al.*, 2014). The mixture is said to be able to increase the medicinal properties of individual species, reduce toxicity and improve the taste acceptance (Guimaraes *et al.*, 2013). In traditional chinese medicine (TCM), some of the anti-dementia herbal formulation have been proved scientifically to exert dementia inhibition effect (Kong *et al.*, 2009). The different herbs exert different effects and when combined, it acts either in synergistic, antagonistic or additive manner. The knowledge regarding the herbal mixture have been passed from older generation to the current generation of herbal practitioner and skillfull healer. Traditionally, the herbal plants that was prepared as mixture have similar therapeutic effects and benefits with possible of synergistic effects (Boer & Cotingting, 2014; Guimaraes *et al.*, 2013). Based on the documentation by Taylor & Wong (1987) in Jakun community, some ailments could be treated by mixture of medicinal plants. Thus, the documentation of medicinal plants for women's healthcare maintenance

and investigation on the herbal mixture used by Jakun women for anti-aging might provide the baseline data for the exploration of anti-aging treatment.

1.2 Problem statement

Aging has been associated with undesirable effects towards physiology and psychology of a person. Although many treatments have been applied to cure or delay these pathologies, it have been found to be ineffective or associated with side effects. Nutricosmetic product with anti-aging claim has increased its popularity in the market, eventhough the efficacy of this product has not been proven scientifically (Draelos, 2010).

Furthermore, the prevalence of neurodegenerative diseases also increase yearly. According to Alzheimer's Disease Foundation Malaysia (ADFM), currently, it is estimated that 50,000 Malaysians suffer from Alzheimer's disease. The incidence of AD could double every five years beyond the age of 65. Although some of the drugs been approved by United State Food and Drug Administration (FDA) for the treatment of AD (i.e Donepezil, Galantamine, Rivastigmine, and Huperzine A), the outcomes are often unsatisfactory (Natarajan *et al.*, 2013; Association, 2014) and associated with side effects such as headache, diarrhoea, drowsiness and vomiting among others.

With regards to the mentioned problems, there is a great demand to develop a safe and more effective cure for skin and brain aging, and medicine to cure and treat these undesirable conditions. Traditional knowledge which is one of the resources in drug discovery is vulnerable to multiple factors, such as erosion and modernization (WIPO, 2001). Thus, the documentation of TK and the investigation on the bioactivities of selected plants might preserve the knowledge and validate the traditional claim.

1.3 Objectives

This study aimed to integrate the traditional knowledge of indigenous people for women health maintenance and to investigate scientifically the potential of the medicinal plants for anti-aging. This aim could be achieved through following specific objectives:

- i. To document the knowledge of Jakun people on medicinal plants for women healthcare.
- ii. To determine the optimal formulations of selected herbal mixture used by Jakun women based on phytochemicals content and antioxidant activities.
- iii. To determine the anti-wrinkle (elastase and collagenase inhibition activities) and anti-hyperpigmentation (tyrosinase inhibition activity) of optimal formulations.
- iv. To determine the anti-Alzheimer potential (acetylcholinesterase inhibition and butyrylcholinesterase inhibition activities) of optimal formulations.
- v. To profile the phytochemical constituents of the selected plants extracts that potentially contributed to the anti-aging properties of optimal formulations.

1.4 Significance of study

In recent years, increasing attention has been paid in searching for solution to reduce the effect of aging. While most of the products with anti-aging claim contain synthetic chemicals which harmful to human, medicinal plants with diverse phytochemicals has been shown to be a better solution as anti-aging agent (Kapoor, Dureja, & Chadha, 2009a). Findings from this study provide scientific evidence on the potential of medicinal plants as anti-aging agent by employing the traditional knowledge of Jakun people on medicinal plants. Besides that, this finding also provides the evidence on the synergistic effects of herbal mixture that is usually believed to be able to exert more therapeutic effects (Guimaraes *et al.*, 2011) as compared to the single herbs. Thus, the evidence of this study will be useful in selecting the formulations with more health benefits and anti-aging effects.

CHAPTER 2

LITERATURE REVIEW

2.1 Documentation of traditional knowledge

2.1.1 Definition of terms

The Convention on Biological Diversity (CBD) in Article 8 (j) has described the term traditional knowledge as :

.... the knowledge, innovations and practices of indigenous and local communities around the world, developed from experience gained over the centuries and adapted to the local culture and environment, and transmitted orally from generation to generation. It tends to be collectively owned and takes several forms from stories, songs, folklore, proverbs, cultural values, beliefs, rituals, community laws, local language and agricultural practices, including the development of plant species and animal breeds. (MoNRE, 2012)

The term “traditional” does not refer to “old”, but it is rather the traditions that have been passed from generations to generations. It relates to the way the knowledge has been created, preserved and disseminated (Finetti, 2011). Traditional Knowledge (TK) comprise of tangible and non-tangible elements (WIPO, 2012). The tangible element of TK is the genetic resources, while the intangible element is the knowledge. The TK sometimes refers as indigenous knowledge (IK). The IK is actually a subset of TK, where the users are communities, people and nation that are indigenous (Van Overwalle, 2005) and the knowledge is only known by the specific community (Finetti, 2011). The traditional knowledge could be further classified into three categories, which are traditional medicinal, agricultural and ecological knowledge (Van Overwalle, 2005). Traditional medicine is defined as the compilation of knowledge, skills and practices based on theories, belief and

experiences native to different cultures, where understandable or not, used for health maintenance as well as in the prevention, diagnosis, improvement or treatment of mental and physical illness (WHO, 2000)

Documentation of TK is defined as a process of identifying, collecting, organizing, registering or recording TK, in order to maintain, manage, use, disseminate and/or protect TK according to specific aims. This knowledge is not limited to simple photographing, or isolated record of tradition, or written notes, but the isolated act need to undergo comprehensive, thought-through process to be regarded as 'documentation'. The most significant thing that needs to be considered while documenting TK is consultation (with and among indigenous or local people), participation and prior informed consent (PIC) before the documenting process starts (WIPO, 2012).

The traditional medicine usually involves all organisms such as animals, plants, and microbes in the treatment of illness by local or indigenous people. Another term that is usually being used in traditional knowledge field is ethnobotany. Ethnobotany and ethnopharmacology are interdisciplinary field that focused on empirical knowledge of indigenous people regarding medicinal substances, their potential health benefits and side effect associated with such remedies (Gurib-fakim, 2006). Ethnobotany has not only restricted to plants, but also involves the studies of algae, lichens and fungi (Eldeen *et al.*, 2016). In the current study, the utilization of medicinal plants, including fungi were investigated among Jakun women in Kampung Peta.

2.1.2 The importance of documenting the traditional knowledge

2.1.2.1 Conserving genetic resources and natural heritage

The traditional knowledge on the uses of organisms including animals in traditional medical system, is important to be documented as the biodiversity is eroding from day to day (Alves & Rosa, 2007). Destructive exploitation of the tropical forest such as illegal logging and conversion of forest to plantation have become the key to the loss of biodiversity and global climate. The precious medicinal plants which have the potential to be developed as useful drugs might be lost along with the forest degradation. In Malaysia, the TK documentation is increasing. Likewise, Forest Research Institute Malaysia (FRIM) with other organizations and universities are

REFERENCES

- Abu Bakar, M. F., Ismail, N. A., Isha, A., & Ling, A. L. M. (2016). Phytochemical Composition and Biological Activities of Selected Wild Berries (*Rubus moluccanus* L., *R. fraxinifolius* Poir., and *R. alpestris* Blume). *Evidence-Based Complementary and Alternative Medicine*, 10 pages.
- Adnan, N., & Othman, N. (2012). The Relationship Between Plants and The Malay Culture. *Procedia - Social and Behavioral Sciences*, 42, 231–241.
- Ahmad, F. (2000). Chemical Studies on Biologically Active Natural Product from Medicinal Plants in Sabah. *Borneo Science*, 7, 55–62.
- Ahn, S., Halake, K., & Lee, J. (2017). Antioxidant and Ion-Induced Gelation Functions of Pectins Enabled by Polyphenol Conjugation. *International Journal of Biological Macromolecules*, 31(101), 776–782.
- Almeida, A. A. De, & Scarminio, I. S. (2007). Statistical Mixture Design Optimization of Extraction Media and Mobile Phase Compositions for the Characterization of Green Tea. *Journal of Separation Science*, 30, 414–420.
- Alves, R. R. N., & Rosa, I. M. L. (2007). Biodiversity, Traditional Medicine and Public Health: Where Do They Meet? *Journal of Ethnobiology and Ethnomedicine*, 3, 9 pages.
- Amarasingham, R. D., Bisset, N. G., Millard, A. H., & Woods, M. C. (1964). A Phytochemical Survey of Malaya Part III . Alkaloids and Saponins. *Economic Botany*, 18(3), 270–278.
- Analyst, G. I. (2015). MCP-6498 : Nutricosmetics - A Global Strategic Bussiness Report. Retrieved March 6, 2017, from <http://www.strategyr.com/pressMCP-6498.asp>
- Anunciato, T. P., & Filho, P. A. da R. (2012). Carotenoids and Polyphenols in Nutricosmetics, Nutraceuticals, And Cosmeceuticals. *Journal of Cosmetic Dermatology*, 11, 51–54.
- Arung, E. T., Kusuma, I. W., Christy, E. O., Shimizu, K., & Kondo, R. (2009). Evaluation of Medicinal Plants from Central Kalimantan for Antimelanogenesis. *Journal of Natural Medicines*, 63(4), 473–480.

- Aruoma, O. I. (2003). Methodological Considerations for Characterizing Potential Antioxidant Actions of Bioactive Components in Plant Foods. *Fundamental and Molecular Mechanisms of Mutagenesis*, 524, 9–20.
- Ashgar, M. N., Khan, I. U., Zia, I., Ahmad, M., & Qureshi, F. A. (2008). Modified 2,2'-azinobis(3-ethylbenzothiazoline)-6-sulphonic Acid Radical Cation Decolorization Assay for Antioxidant Activity of Human Plasma and Extracts of Traditional Medicinal Plants. *Acta Chimica Slovenica*, 55, 408–418.
- Association, A. (2014). 2014 Alzheimer's Disease Facts and Figures. *Alzheimer's & Dementia*, 10(2), 75 pages.
- Atoui, A. K., Mansouri, A., & Boskou, G. (2005). Tea and Herbal Infusions : Their Antioxidant Activity and Phenolic Profile. *Food Chemistry*, 89, 27–36.
- Atta-ur-Rahman, Choudhary, M. I., & Thomsen, W. J. (2005). *Bioassay Techniques for Drug Development*. Netherland : Hardwood Academic Publisher.
- Atta-ur-Rahman, Parveen, S., Khalid, A., Farooq, A., & Choudhary, M. I. (2001). Acetyl and Butyrylcholinesterase Inhibiting Triterpenoid Alkaloids from *Buxus papillosa*. *Phytochemistry*, 58(6), 963–968.
- Azliza, M. A., Ong, H. C., Vikineswary, S., Noorlidah, A., & Haron, N. W. (2012). Ethno-medicinal resources used by the Temuan in Ulu Kuang village. *Studies on Ethno-Medicine*, 6(1), 17–22.
- Bastianetto, S., & Quirion, R. (2008). Phytochemicals and the Aging Brain. In M. S. Meskin, W. R. Bidlack, & R. K. Randolph (Eds.), *Phytochemicals : Aging and Health*. CRC Press. pp. 107–117
- Baumann, L. S. (2007). Less-known Botanical Cosmeceuticals. *Dermatologic Therapy*, 20, 330–342.
- Baurin, N., Arnoult, E., Scior, T., Do, Q. T., & Bernard, P. (2002). Preliminary Screening of Some Tropical Plants for Anti-tyrosinase Activity. *Journal of Ethnopharmacology*, 82(2–3), 155–158.
- Benzie, I. F. F., & Strain, J. J. (1996). The Ferric Reducing Ability of Plasma (FRAP) as a Measure of “Antioxidant Power”: The FRAP Assay. *Analytical Biochemistry*, 239(1), 70–76.
- Bochi, V. C., Barcia, M. T., Rodrigues, D., Speroni, C. S., Giusti, M. M., & Godoy, H. T. (2015). Polyphenol Extraction Optimisation from Ceylon gooseberry (*Dovyalis hebecarpa*) Pulp. *Food Chemistry*, 164(2014), 347–354.

- Boer, H. J. De, & Cotingting, C. (2014). Medicinal Plants for Women's Healthcare in Southeast Asia: A Meta-Analysis of Their Traditional Use, Chemical Constituents, and Pharmacology. *Journal of Ethnopharmacology*, 151(2), 747–767.
- Boğa, M., Alkan, H., Ertaş, A., Oral, E. V., Yılmaz, A., Yeşil, Y., Gören, A.C., Temel, H., & Kolak, U. (2016). Phytochemical Profile and Some Biological Activities of Three *Centaurea* species from Turkey. *Tropical Journal of Pharmaceutical Research*, 15(9), 1865–1875.
- Bolognesi, M. L., Matera, R., Minarini, A., Rosini, M., & Melchiorre, C. (2009). Alzheimer's Disease: New Approaches to Drug Discovery. *Current Opinion in Chemical Biology*, 13, 303–308.
- Brand-Williams, W., Cuvelier, M. E., & Berset, C. (1995). Use of a Free Radical Method to Evaluate Antioxidant Activity. *LWT - Food Science and Technology*, 28(1), 25–30.
- Camera, E., Mastrofrancesco, A., Fabbri, C., Daubrawa, F., Picardo, M., Sies, H., & Stahl, W. (2009). Astaxanthin, Canthaxanthin and Beta-Carotene Differently affect UVA-Induced Oxidative Damage and Expression of Oxidative Stress-Responsive Enzymes. *Experimental Dermatology*, 18, 222–31.
- Capitani, C. D., Carvalho, A. C. L., Botelho, P. B., Carrapeiro, M. M., & Castro, I. A. (2009). Synergism on Antioxidant Activity between Natural Compounds Optimized by Response Surface Methodology. *European Journal of Lipid Science and Technology*, 111, 1100–1110.
- Carabajal, M. P. A., Isla, M. I., & Zampini, I. C. (2016). Evaluation of Antioxidant and Antimutagenic Activity of Herbal Teas from Native Plants used in traditional Medicine in Argentina. *South African Journal of Botany*, 110, 258–265.
- Carocho, M., & Ferreira, I. C. F. R. (2013). A Review on Antioxidants, Prooxidants and Related Controversy: Natural and Synthetic Compounds, Screening and analysis Methodologies and Future Perspectives. *Food and Chemical Toxicology*, 51(1), 15–25.
- CBD, S. (2002). *Bonn guidelines on access to genetic resources and fair and equitable sharing of the benefits arising out of their utilization*. Montreal: Secretariat of Convention on Biological Diversity.

- Chang, L. W., Juang, L. J., Wang, B. Sen, Wang, M. Y., Tai, H. M., Hung, W. J., Chen, Y. J., & Huang, M. H. (2011). Antioxidant and Antityrosinase Activity of Mulberry (*Morus alba* L.) twigs and root bark. *Food and Chemical Toxicology*, 49(4), 785–790.
- Chang, T. S. (2009). An Updated Review of Tyrosinase Inhibitors. *International Journal of Molecular Sciences*, 10(6), 2440–2475.
- Chang, Y. S., & Lee, S. S. (2004). Utilisation of Macrofungi Species in Malaysia. *Fungal Diversity*, 15–22.
- Chemat, F., Vian, M. A., & Cravotto, G. (2012). Green Extraction of Natural Products : Concept and Principles. *International Journal of Molecular Sciences*, 13, 8615–8627.
- Cheung, D. W., Koon, C., Ng, C., Leung, P., Fung, K.-P., Poon, S. K.-S., & Lau, C. B.-S. (2012). The roots of *Salvia miltiorrhiza* (Danshen) and *Pueraria lobata* (Gegen) inhibit atherogenic events : A study of the combination effects of the 2-herb formula. *Journal of Ethnopharmacology*, 143, 859–866.
- Chompoo, J., Upadhyay, A., Fukuta, M., & Tawata, S. (2012). Effect of *Alpinia zerumbet* components on antioxidant and skin diseases-related enzymes. *BMC Complementary and Alternative Medicine*, 12, 106.
- Dej-adisai, S., Tinpun, K., Wattanapiromsakul, C., & Keawpradub, N. (2015). Bio-activities and Phytochemical Investigation of *Cnestis palala* (Lour.) Merr. *African Journal of Traditional, Complementary and Alternative Medicine*, 12, 27–37.
- Dey, A., Bhattacharya, R., Mukherjee, A., & Kumar, D. (2017). Natural products against Alzheimer's Disease : Pharmaco-therapeutics and Biotechnological Interventions. *Biotechnology Advances*, 35(2), 178–216.
- Draelos, Z. D. (2010). Nutrition and Enhancing Youthful-Appearing Skin. *Clinics in Dermatology*, 28(4), 400–408.
- Dua, D., & Srivastava, N. S. (2016). A study on antioxidant and anti-aging properties of few medicinal plants. *International Journal of Pharmacy and Pharmaceutical Science*, 8(1), 344–347.
- Duque, L., Bravo, K., & Osorio, E. (2017). A Holistic Anti-Aging Approach Applied in Selected Cultivated Medicinal Plants: A View of Photoprotection of The Skin by Different Mechanisms. *Industrial Crops and Products*, 97, 431–439.

- Dzialo, M., Mierziak, J., Korzun, U., Preisner, M., Szopa, J., & Kulma, A. (2016). The Potential of Plant Phenolics in Prevention and Therapy of Skin Disorders. *International Journal of Molecular Sciences*, 17 (160), 1-40.
- Eldeen, I. M. S., Effendy, M. A. W., & Muhammad, T. S. T. (2016). Ethnobotany : Challenges and future perspectives. *Research Journal of Medicinal Plants*, 10 (6), 382–387.
- Ellman, G. L., Courtney, K. D., Jr., V. A., & Featherstone, R. M. (1961). A New and Rapid Colorimetric Determination of Acetylcholinesterase Activity. *Biochemical Pharmacology*, 7, 88–95.
- Embuscado, M. E. (2015). Spices and Herbs: Natural Sources of Antioxidants – A Mini Review. *Journal of Functional Foods*, 18, 811–819.
- Fadaeinasab, M., Basiri, A., Kia, Y., Ali, H. M., Karimian, H., & Murugaiyah, V. (2015). New Indole Alkaloids from the Bark of *Rauvolfia reflexa* and their cholinesterase inhibitory activity. *Cell Physiology and Biochemistry*, 37, 1997–2011.
- Farzaneh, V., & Carvalho, I. S. (2015). A Review of the Health Benefit Potentials of Herbal Plant Infusions and their Mechanism of Actions. *Industrial Crops and Products*, 65, 247–258.
- Fernando, C. D., & Soysa, P. (2015). Extraction Kinetics of Phytochemicals and Antioxidant Activity during Black Tea (*Camellia sinensis* L.) Brewing. *Nutrition Journal*, 14 (1), 74.
- Finetti, C. (2011). Traditional Knowledge and the Patent System : Two Worlds Apart?. *World Patent Information*, 33 (1), 58–66.
- Floegel, A., Kim, D. O., Chung, S. J., Koo, S. I., & Chun, O. K. (2011). Comparison of ABTS/DPPH Assays to Measure Antioxidant Capacity in Popular Antioxidant-Rich US Foods. *Journal of Food Composition and Analysis*, 24(7), 1043–1048.
- Fui, L. H., Haron, N., Yusof, N., Baharuddin, I. N., Mustapha, N. M., Tan, A. L., Zaidi, N. H., Jamaludin, F., & Mohtar, M. (2015). Documenting Traditional Forest-Related Knowledge of Medicinal Plants in Malaysia : A Comprehensive Approach. *Journal of Tropical Resources and Sustainable Science*, 3, 77–85.
- German-Baez, L. J., Valdez-Flores, M., Figueroa-Perez, M. G., Garduno-Felix, K. G., Valdez-Ortiz, R., Meza-Ayala, K. A., & Valdez-Ortiz, A. (2017). Anti-aging and Nutraceutical Characterization of Plant Infusions used in Traditional

- Medicine. *Pakistan Journal of Nutrition*, 16 (4), 285–292.
- Ghimeray, A. K., Jung, U. S., Lee, H. Y., Kim, Y. H., Ryu, E. K., & Chang, M. S. (2015). *In Vitro* Antioxidant, Collagenase Inhibition, and *in vivo* Anti-wrinkle Effects of Combined Formulation Containing *Punica granatum*, *Ginkgo biloba*, *Ficus carica*, and *Morus alba* fruits extract. *Clinical, Cosmetic and Investigational Dermatology*, 8, 389–396.
- Goncalves, S., Gomes, D., Costa, P., & Romano, A. (2013). The Phenolic Content and Antioxidant Activity of Infusions from Mediterranean Medicinal Plants. *Industrial Crops and Products*, 43, 465–471.
- Good, C. M. (1987). *Ethnomedical Systems in Africa : Patterns of Traditional Medicine in Rural and Urban Kenya*. Hove: Guilford Press.
- Granato, D., Grevink, R., Zielinski, A. F., Nunes, D. S., & Ruth, S. M. Van. (2014). Analytical Strategy Coupled with Response Surface Methodology to Maximize the Extraction of Antioxidants from Ternary Mixtures Of Green, Yellow, and Red Teas (*Camellia sinensis* var. *sinensis*). *Journal of Agricultural and Food Chemistry*, 62, 10283–10296.
- Granato, D., Prado-silva, L. do, Alvarenga, V. O., Zielinski, A. A. F., Bataglion, G. A., Morais, D. R. de, Eberlin, M. N. & Sant'Ana, A. de S. (2016). Characterization of Binary and Ternary Mixtures of Green, White and Black Tea Extracts by Electrospray Ionization Mass Spectrometry and Modeling of their *In Vitro* Antibacterial Activity. *LWT - Food Science and Technology*, 65, 414–420.
- Guimaraes, R., Barros, L., Carvalho, A. M., & Ferreira, I. C. F. R. (2011). Infusions and Decoctions of Mixed Herbs used in Folk Medicine: Synergism in Antioxidant Potential. *Phytotherapy Research*, 25(8), 1209–1214.
- Guimaraes, R., Barros, L., Duenas, M., Calhelha, R. C., Carvalho, A. M., & Santos-Buelga, C., Ferreira, I. C. F. R. (2013). Infusion and Decoction of Wild German Chamomile: Bioactivity and Characterization Of Organic Acids And Phenolic Compounds. *Food Chemistry*, 136 (2), 947–954.
- Gurib-fakim, A. (2006). Medicinal plants : Traditions of yesterday and drugs of tomorrow. *Molecular Aspects of Medicine*, 27, 1–93.
- Handa, C. L., De Lima, F. S., Guelfi, M. F. G., Georgetti, S. R., & Ida, E. I. (2016). Multi-response Optimisation of the Extraction Solvent System for Phenolics and Antioxidant Activities from Fermented Soy Flour using a Simplex-Centroid

- Design. *Food Chemistry*, 197, 175–184.
- Harman, D. (1956). Aging: A Theory Based on Free Radical and Radiation Chemistry. *Journal of Gerontology*, 11 (3), 298–300.
- Harman, D. (1998). Free Radical Theory of Ageing: Applications. *Free Radicals and Aging*, 7 (3), 169–177.
- Hayflick, L. (1965). The Limited *In Vitro* Lifetime of Human Diploid Cell Strains. *Experimental Cell Research*, 37 (3), 614–636.
- Heinrich, M., & Teoh, H. L. (2004). Galanthamine from Snowdrop — The Development of a Modern Drug against Alzheimer's Disease from Local Caucasian Knowledge. *Journal of Ethnopharmacology*, 92, 147–162.
- Ho, Y., So, K., & Chang, R. C. (2010). Anti-aging Herbal Medicine - How and Why Can they be used in Aging-associated Neurodegenerative Diseases? *Ageing Research Reviews*, 9(3), 354–362.
- Hong, Y.-H., Jung, E. Y., Noh, D. O., & Suh, H. J. (2014). Physiological Effects of Formulation containing Tannase-Converted Green Tea Extract On Skin Care: Physical Stability, Collagenase, Elastase, and Tyrosinase Activities. *Integrative Medicine Research*, 3(1), 25–33.
- Huang, D., & Prior, R. L. (2005). The Chemistry behind Antioxidant Capacity Assays. *Journal of Agricultural and Food Chemistry*, 53, 1841–1856.
- Irrera, N., Pizzino, G., Anna, R. D., Vaccaro, M., Arcoraci, V., Squadrito, F., Altavilla, D., & Bitto, A. (2017). Dietary management of skin health: The role of genistein. *Nutrients*, 9 (622), 1–10.
- Irshad, M., & Chaudhuri, P. S. (2002). Oxidant-antioxidant System: Role and Significance in Human Body. *Indian Journal of Experimental Biology*, 40, 1233–1239.
- Ishola, I. O., Tota, S., Adeyemi, O. O., Agbaje, E. O., Narender, T., & Shukla, R. (2013). Protective Effect of *Cnestis ferruginea* and its Active Constituent on Scopolamine-Induced Memory Impairment in Mice: A Behavioral and Biochemical Study. *Pharmaceutical Biology*, 51(7), 825–835.
- Ismail, I., Linatoc, A. C., Mohamed, M., & Tokiman, L. (2015). Documentation of Medicinal Plants Traditionally used by the Jakun people of Endau-Rompin (PETA) for Treatments of Malaria-like Symptoms. *Jurnal Teknologi*, 77(31), 63–69.

- Jacobo-Velazquez, D. A., & Cisneros-Zevallos, L. (2009). Correlations of antioxidant activity against phenolic content revisited: A new approach in data analysis for food and medicinal plants. *Journal of Food Science*, 74(9), 107–113.
- JAKOA. (2017). Tribes/Nation. Retrieved February 7, 2017, from <http://www.jakoa.gov.my/en/orang-asli/info-orang-asli/suku-kaumbangsa/>
- Janser, R., Castro, S. De, & Sato, H. H. (2013). Comparison and Synergistic Effects of Intact Proteins and their Hydrolysates On The Functional Properties and Antioxidant Activities in a Simultaneous Process of Enzymatic Hydrolysis. *Food and Bioproducts Processing*, 92 (1), 80–88.
- Janser, R., Castro, S. De, Sato, H. H., De Castro, R. J. S., & Sato, H. H. (2013). Synergistic Effects of Agroindustrial Wastes on Simultaneous Production of Protease and α -Amylase under Solid State Fermentation using A Simplex Centroid Mixture Design. *Industrial Crops and Products*, 49, 813–821.
- Jin, C. B. (2005). Medicinal Properties and Common Usages of Some Palm Species in the Kampung Peta Community of Endau-Rompin National Park, Johor. *Journal of Tropical Medicinal Plants*, 6(I), 79–83.
- Kamagaju, L., Morandini, R., Bizuru, E., Nyetera, P., Nduwayezu, J. B., Stévigny, C., Ghanem, G., & Duez, P. (2013). Tyrosinase Modulation by Five Rwandese Herbal Medicines Traditionally used for Skin Treatment. *Journal of Ethnopharmacology*, 146 (3), 824–834.
- Kang, T. H., Park, H. M., Kim, Y. B., Kim, H., Kim, N., Do, J. H., Kang, C., Cho, Y., & Kim, S. Y. (2009). Effects of red ginseng extract on UVB irradiation-induced skin aging in hairless mice. *Journal of Ethnopharmacology*, 123 (3), 446–451.
- Kankara, S. S., Ibrahim, M. H., Mustafa, M., & Go, R. (2015). Ethnobotanical Survey of Medicinal Plants used for Traditional Maternal Healthcare in Katsina State, Nigeria. *South African Journal of Botany*, 97, 165–175.
- Kapoor, V. K., Dureja, J., & Chadha, R. (2009a). Herbals in the Control of Ageing. *Drug Discovery Today*, 14 (19/20), 992–998.
- Kapoor, V. K., Dureja, J., & Chadha, R. (2009b). Synthetic Drugs with Anti-Ageing Effects. *Drug Discovery Today*, 14 (17/18), 899–904.
- Kardooni, R., Kari, F. B., Rohani, S., Yahaya, B., & Yusup, S. H. (2014). Traditional Knowledge of Orang Asli on Forests in Peninsular Malaysia. *Indian Journal of Traditional Knowledge*, 13, 283–291.

- Karim, A. A., Azlan, A., Ismail, A., Hashim, P., Abd Gani, S. S., Zainudin, B. H., & Abdullah, N. A. (2014). Phenolic Composition, Antioxidant, Anti-wrinkles and Tyrosinase Inhibitory Activities of Cocoa Pod Extract. *BMC Complementary and Alternative Medicine*, 14, 381.
- Karimi, E., Jaafar, H. Z. E., & Ahmad, S. (2011). Phytochemical Analysis and Antimicrobial Activities of Methanolic Extracts of Leaf, Stem and Root from Different Varieties Of *Labisa pumila* Benth. *Molecules*, 16 (6), 4438–4450.
- Khan, B. A., Akhtar, N., Hussain, I., Abbas, K. A., & Rasul, A. (2013). Whitening efficacy of Plant Extracts including *Hippophae rhamnoides* and *Cassia fistula* Extracts on the Skin of Asian Patients with Melasma. *Postepy Dermatologii I Alergologii*, 4, 226–233.
- Khazanah Endau Rompin Herba*. (2007) (1st ed.). Kuala Lumpur: Utusan Publication & Distributors Sdn. Bhd.
- Khazanah Endau Rompin Herba*. (2008) (2nd ed.). Skudai: Perbadanan Bioteknologi dan Biodiversiti Negeri Johor.
- Khoddami, A., Wilkes, M. A., & Roberts, T. H. (2013). Techniques for Analysis of Plant Phenolic Compounds. *Molecules*, 18 (2), 2328–2375.
- Khor, G. L., & Shariff, Z. M. (2008). The Ecology of Health and Nutrition of “Orang Asli” (Indigenous people) Women and Children in Peninsular Malaysia. *Tribes and Tribals*, 2, 67–77.
- Kim, D. B., Shin, G. H., Kim, J. M., Kim, Y. H., Lee, J. H., Lee, J. S., Seng, H. J., Choe, S. Y., Park, I. J., Cho, J. H., & Lee, O. H. (2016). Antioxidant and Anti-ageing Activities of Citrus-based Juice Mixture. *Food Chemistry*, 194, 920–927.
- Kim, H. M., An, H. S., Bae, J.-S., Kim, J. Y., Choi, C. H., Kim, J. Y., Lim, J. H., Choi, J.-H., Song, H., Moon, S. H., Park, Y. J., Chang, S.-J., & Choi, S. Y. (2017). Effects of Palmitoyl-KVK-L-Ascorbic Acid on Skin Wrinkles and Pigmentation. *Archives of Dermatological Research*, 309 (5): 397-402.
- Kim, M., Ko, J., & Lim, S. (2016). Formulation Optimization of Antioxidant-Rich Juice Powders based on Experimental Mixture Design. *Journal of Food Processing and Preservation*, 10 pages.
- Kim, Y. H., Chung, C. B., Kim, J. G., Ko, K. I. K., Park, S. H., Kim, J., Kim, J. H., Eom, S. Y., Kim, Y. S., Hwang, Y. I. & Kim, K. H. (2008). Anti-wrinkle Activity of Ziyuglycoside I isolated from a *Sanguisorba officinalis* Root Extract and its Application as a Cosmeceutical Ingredient. *Bioscience, Biotechnology*,

and *Biochemistry*, 72(2), 303–311.

- Kim, Y. J., & Uyama, H. (2005). Tyrosinase Inhibitors from Natural and Synthetic Sources: Structure, Inhibition Mechanism and Perspective for the Future. *Cellular and Molecular Life Sciences*, 62 (15), 1707–1723.
- Kong, D.-X., Li, X.-J., & Zhang, H.-Y. (2009). Where is the Hope for Drug Discovery? Let History Tell the Future. *Drug Discovery Today*, 14(3/4), 115–119.
- Kyung, Y., Soo, K., Choi, J., Rong, C., Kyeom, J., Kim, K. Y., Choi, J. H., Song, S. W., Kim, C. J., Park, G. G., Park, C. S., & Shin, C. P. D. (2016). Antioxidant and Cognitive-Enhancing Activities of *Arctium lappa* L. roots in AB1-42-induced Mouse Model. *Applied Biological Chemistry*, 59 (4), 553–565.
- Labat-Robert, J., Fourtanier, A., Boyer-Lafargue, B., & Robert, L. (2000). Age Dependent Increase of Elastase Type Protease Activity in Mouse Skin Effect of UV-Irradiation. *Journal of Photochemistry and Photobiology B: Biology*, 57, 113–118.
- Lar, M. S. M. (2014). Phytochemical Analysis of *Cnestis palala* (Lour.) Merr. and its Antimicrobial Activity. *Universities Research Journal*, 6 (1), 161–172.
- Lawless, L. J. R., Threlfall, R. T., Meullenet, J. F., & Howard, L. R. (2013). Applying a Mixture Design for Consumer Optimization of Black Cherry, Concord Grape and Pomegranate Juice Blends. *Journal of Sensory Studies*, 28 (2), 102–112.
- Lee, K.-K., Kim, J.-H., Cho, J.-J., & Choi, J.-D. (1999). Inhibitory Effects Of 150 Plant Extracts on Elastase Activity, and their Anti-Inflammatory. *International Journal of Cosmetic Science*, 21, 71–82.
- Lee, K. O., Kim, S. N., & Kim, Y. C. (2014). Anti-wrinkle Effects of Water Extracts of Teas in Hairless Mouse. *Toxicological Research*, 30(4), 283–289.
- Lee, S.-H., & Min, K.-J. (2017). Lifespan-Extending Effect of Resveratrol and Other Phytochemicals. In Alexander M. Vaiseman (Ed.). *Practice, Anti-aging Drugs : From Basic Research to Clinical*. United Kingdom: The Royal Society of Chemistry. pp. 328–351.
- Lee, S. J., Koh, J., Ha, B., & Boo, Y. C. (2010). *Quercus glauca* Extract and Rutin inhibit the UVB-induced Expression of Matrix Metalloproteinase-1 in Human Dermal Fibroblasts. *Journal of the Korean Society for Applied Biological Chemistry*, 53(6), 677–684.

- Lee, S., Oh, D., Lee, S., Kim, G. R., Lee, J. S., Son, Y. K., Bae, C. H., Yeo, J., & Lee, C. H. (2015). Chemotaxonomic metabolite profiling of 62 indigenous plant species and its correlation with bioactivities. *Molecular Nutrition and Food Research*, 19719–19734.
- Lee, S. S., Chang, Y. S., & Noraswati, M. N. R. (2009). Utilization of Macrofungi by Some Indigenous Communities for Food and Medicine in Peninsular Malaysia. *Forest Ecology and Management*, 257, 2062–2065.
- Level, W. E. E. (2017). Decamethylcyclopentasiloxane (D5). *Toxicology and Industrial Health*, 33 (1), 16–27.
- Li, S., Li, S. K., Li, H. Bin, Xu, X. R., Deng, G. F., & Xu, D. P. (2014). Antioxidant Capacities of Herbal Infusions. In Preedy, V. (Ed.). *Processing and Impact on Antioxidants in Beverages*. United Kingdom : Academic Press. pp. 41-50
- Limtrakul, P., Yodkeeree, S., Thippraphan, P., Punfa, W., & Srisomboon, J. (2016). Anti-aging and Tyrosinase Inhibition Effects of *Cassia fistula* Flower Butanolic Extract. *BMC Complementary and Alternative Medicine*, 16 (1), 497.
- Lin, K. W. (2005). Ethnobotanical Study of Medicinal Plants used by Jah Hut Peoples in Malaysia. *Indian Journal of Medical Sciences*, 59 (4), 156–161.
- Lobo, V., Patil, A., Phatak, A., & Chandra, N. (2010). Free Radicals, Antioxidants and Functional Foods : Impact on Human Health. *Pharmacognosy Review*, 4 (8), 118–126.
- Lusia Barek, M., Hasmadi, M., Zaleha, A. Z., & Mohd Fadzelly, A. B., (2015). Effect of Different Drying Methods on Phytochemicals and Antioxidant Properties of Unfermented and Fermented Teas from Sabah Snake Grass (*Clinacanthus nutans* Lind.) leaves. *International Food Research Journal*, 22(2), 661–670.
- Mandrone, M., Lorenzi, B., Venditti, A., Guarcini, L., Bianco, A., Sanna, C., Ballero, M., Poli, F., & Antognoni, F. (2015). Antioxidant and Anti-collagenase Activity of *Hypericum hircinum* L. *Industrial Crops and Products*, 76, 402–408.
- Martin, G. J. (1995). *Ethnobotany : A Method Manual*. London: Chapman & Hall.
- Masaki, H. (2010). Role of Antioxidants in the Skin: Anti-aging Effects. *Journal of Dermatological Science*, 58 (2), 85–90.
- Masamoto, Y., Lida, S., & Kubo, M. (1980). Inhibitory Effect of Chinese Crude Drugs on Tyrosinase. *Journal of Medicinal Plant Research*, 40 (620), 361–365.

- Masron, T., Masami, F., & Ismail, N. (2013). Orang Asli in Peninsular Malaysia : Population, Spatial Distribution and Socio-Economic Condition. *Journal of Ritsumeikan Social Sciences and Humanities*, 6, 75–115.
- Masuda, M., Murata, K., Fukuham, A., Naruto, S., Fujita, T., Uwaya, A. & Matsuda, H. (2009). Inhibitory Effects of Constituents of *Morinda citrifolia* seeds on Elastase and Tyrosinase. *Journal of Natural Medicines*, 63(3), 267–273.
- Matés, J. M., Pérez-Gómez, C., & Núñez de Castro, I. (1999). Antioxidant Enzymes and Human Diseases. *Clinical Biochemistry*, 32 (8), 595–603.
- Mathew, M., & Subramanian, S. (2014). *In Vitro* Screening for Anti-cholinesterase and Antioxidant Activity of Methanolic Extracts of Ayurvedic Medicinal Plants used for Cognitive Disorders. *PLOS ONE*, 9 (1), 1–7.
- Mazlan, N. A., Mediani, A., Abas, F., Ahmad, S., Shaari, K., Khamis, S. & Lajis, N. H. (2013). Antioxidant, Antityrosinase, Anticholinesterase, and Nitric Oxide Inhibition Activities of Three Malaysian *Macaranga* Species. *The Scientific World Journal*, 8 pages.
- McClain, C. (1989). *Women as healers: Cross-cultural perspectives*. New Brunswick and London: University Press.
- Mensor, L. L., Menezes, F. S., Leitao, G. G., Reis, A. S., Santos, T. C. dos, Coube, C. S., & Leitao, S. G. (2001). Screening of Brazilian Plant Extracts for Antioxidant Activity by the use of DPPH Free Radical Method. *Phytotherapy Research*, 15, 127–130.
- Merzouki, B., Ed-Derfoufi, F., & Mesa, J. F. (2000). Contribution to the Knowledge of Rifian Traditional Medicine. II: Folk medicine in Ksar Lakbir district (NW Marocco). *Fitoterapia*, 71, 178–307.
- Mohammad, N. S., Milow, P., & Ong, H. C. (2012a). Traditional Knowledge on Medicinal Plant among the Orang Asli Villagers in Kampung Lubuk Ulu Legong, Kedah, Malaysia. In *Technology, Science, Social Sciences and Humanities International Conference 2012*. 9 pages.
- Mohammad, N. S., Milow, P., & Ong, H. C. (2012b). Traditional Medicinal Plants Used by The Kensiu Tribe of Lubuk. *Studies on Ethno-Medicine*, 6 (3), 149–153.
- Momtaz, S., Mapunya, B. M., Houghton, P. J., Edgerly, C., Hussein, A., Naidoo, S., & Lall, N. (2008). Tyrosinase Inhibition by Extracts And Constituents of *Sideroxylon inerme* L. Stem Bark, used in South Africa for Skin Lightening.

Journal of Ethnopharmacology, 119(3), 507–512.

MoNRE (2012). *National Policy on Biological Diversity 2016-2025*. Putrajaya : Ministry of Natural Resources and Environment.

Muddathir, A. M., Yamauchi, K., Batubara, I., Mohieldin, E. A. M., & Mitsunaga, T. (2017). Anti-tyrosinase, Total Phenolic Content and Antioxidant Activity of Selected Sudanese Medicinal Plants. *South African Journal of Botany*, 109, 9–15.

Mukherjee, P. K., Kumar, V., Mal, M., & Houghton, P. J. (2007). Acetylcholinesterase Inhibitors from Plants. *Phytomedicine*, 14 (4), 289–300.

Mukherjee, P. K., Maity, N., Nema, N. K., & Sarkar, B. K. (2011). Bioactive Compounds from Natural Resources against Skin Aging. *Phytomedicine*, 19 (1), 64–73.

Mukherjee, P. K., Maity, N., Nema, N. K., & Sarkar, B. K. (2013). Natural Matrix Metalloproteinase Inhibitors: Leads from Herbal Resources. In *Studies in Natural Products Chemistry*. 1st ed. United Kingdom : Elsevier. pp. 91–113.

Munhoz, V. M., Longhinia, R., Souza, J. R. P., Zequic, J. A. C., Mello, E. V. S. L., Lopes, G. C., & Mello, J. C. P. (2014). Extraction of Flavonoids from *Tagetes patula* : Process Optimization and Screening for Biological Activity. *Revista Brasileira de Farmacognosia*, 24 (5), 576–583.

Mushtaq, A., N Rasool, Riaz, M., Tareen, R. B., Zubair, M., Rashid, U., Khan, M. A., & Taufiq-Yap, Y. H. (2013). Characterization of Essential Oil and Fixed Oil of *Clematis Graveolens* by GC-MS and their Antioxidant, Antimicrobial Studies. *Oxidation Communications*, 36 (4), 1067–1078.

Najmuddin, S., & Rahman, N. A. (2017). Comparative Study of Antioxidant Level and Activity from Leaf Extracts of *Annona muricata* Linn obtained from Different Locations. *Pertanika*, 40 (1), 119–130.

Narayanaswamy, R., & Ismail, I. S. (2015). Cosmetic potential of Southeast Asian herbs: an overview. *Phytochemistry Reviews*, 14 (3), 419–428.

Natarajan, S., Shunmugiah, K. P., & Kasi, P. D. (2013). Plants Traditionally used in Age-Related Brain Disorders (Dementia): An Ethanopharmacological Survey. *Pharmaceutical Biology*, 51 (4), 492–523.

Ndhhlala, A. R., Aremu, A. O., Moyo, M., Amoo, S. O., & Van Staden, J. (2012). Acetylcholinesterase Inhibitors from Plant Sources: Friends or foes?. In White, C. J. & Tait, J. E. *Cholinesterase: Production, Uses and Health Effects*. New

York : Nova Science Publisher. pp. 1-25.

Ndhlala, A. R., Moyo, M., & Van Staden, J. (2010). Natural Antioxidants: Fascinating or Mythical Biomolecules?. *Molecules*, 15(10), 6905–6930.

Ndlovu, G., Fouche, G., Tselanyane, M., Cordier, W., & Steenkamp, V. (2013). *In Vitro* Determination of the Anti-aging Potential of Four Southern African Medicinal Plants. *BMC Complementary and Alternative Medicine*, 13(1), 7 pages.

Neagu, E., Radu, G. L., Albu, C., & Paun, G. (2016). Antioxidant Activity, Acetylcholinesterase and Tyrosinase Inhibitory Potential of *Pulmonaria officinalis* and *Centarium umbellatum* Extracts. *Saudi Journal of Biological Sciences*, *In press*.

Nema, N. K., Maity, N., Sarkar, B. K., & Mukherjee, P. K. (2013). Matrix Metalloproteinase, Hyaluronidase and Elastase Inhibitory Potential of Standardized Extract of *Centella asiatica*. *Pharmaceutical Biology*, 51 (9), 1182–1187.

Nema, N. K., Maity, N., Sarkar, B., & Mukherjee, P. K. (2011). *Cucumis sativus* Fruit-Potential Antioxidant, Anti-Hyaluronidase, and Anti-Elastase Agent. *Archives of Dermatological Research*, 303, 247–252.

Norhaiza, M., Maziah, M., & Hakiman, M. (2009). Antioxidative Properties Of Leaf Extracts of a Popular Malaysian Herb, *Labisia pumila*. *Journal of Medicinal Plants Research*, 3 (4), 217–223.

Nour, A. H., Khan, M., Sulaiman, A. Z., Batool, T., Nour, A. H., Khan, M. M., & Kormin, F. (2014). *In Vitro* Anti Acetylcholinesterase and Antioxidant Activity of Selected Malaysian Plants. *Asian Journal of Pharmaceutical and Clinical Research*, 7(3), 93-97.

Omar, S. H., Scott, C. J., Hamlin, A. S., & Obied, H. K. (2017). The Protective Role of Plant Biophenols in Mechanisms of Alzheimer's Disease. *The Journal of Nutritional Biochemistry*, 47, 1–20.

Ong, H. C., Faezah, A. W., & Milow, P. (2012). Medicinal Plants used by the Jah Hut Orang Asli at Kampung Pos Penderas, Pahang, Malaysia. *Studies on Ethno-Medicine*, 6 (1), 11–15.

Ong, H. C., Lina, E., & Milow, P. (2012). Traditional Knowledge and Usage Of Medicinal Plants among the Semai Orang Asli at Kampung Batu 16, Tapah, Perak, Malaysia. *Studies on Ethno-Medicine*, 6(3), 207–211.

- Ong, H. C. U., & Nordiana, M. (1999). Malay Ethno-Medico Botany in Machang , Kelantan , Malaysia. *Fitoterapia*, 502–513.
- Ong, H. C., Zuki, R. M., & Milow, P. (2011). Traditional Knowledge of Medicinal Plants among the Malay villagers in Kampung Mak Kemas, Terengganu, Malaysia. *Studies on Ethno-Medicine*, 5 (3), 175–185.
- Orhan, I., Kartal, M., Tosun, F., & Bilge, S. (2007). Screening of Various Phenolic Acids and Flavonoid Derivatives for their Anticholinesterase Potential. *Zeitschrift Für Naturforschung C*, 62 (11–12), 829–832.
- Orhan, I., Şener, B., Choudhary, M. I., & Khalid, A. (2004). Acetylcholinesterase and Butyrylcholinesterase Inhibitory Activity of Some Turkish Medicinal Plants. *Journal of Ethnopharmacology*, 91 (1), 57–60.
- Orives, J. R., Galvan, D., Coppo, R. L., Henrique, C., Rodrigues, F., Angilelli, K. G., & Borsato, D. (2014). Multiresponse Optimisation on Biodiesel obtained through a Ternary Mixture of Vegetable Oil and Animal Fat : Simplex-Centroid Mixture Design Application. *Energy Conversion and Management*, 79, 398–404.
- Ouedrhiri, W., Balouiri, M., Bouhdid, S., Moja, S., Chahdi, F. O., Taleb, M., & Greche, H. (2016). Mixture Design of *Origanum compactum*, *Origanum majorana* and *Thymus serpyllum* essential oils : Optimization of their Antibacterial Activities. *Industrial Crops and Products Journal*, 89, 1–9.
- Pandey, K. B., & Rizvi, S. I. (2009). Plant Polyphenols as Dietary Antioxidants in Human Health and Disease. *Oxidative Medicine and Cellular Longevity*, 2 (5), 270–278.
- Panyaphu, K., Van On, T., Sirisa-Ard, P., Srisa-Nga, P., Chansakaow, S., & Nathakarnkitkul, S. (2011). Medicinal plants of the Mien (Yao) in Northern Thailand and their Potential Value in the Primary Healthcare of Postpartum Women. *Journal of Ethnopharmacology*, 135(2), 226–237.
- Patocka, J., Kuca, K., & Jun, D. (2004). Acetylcholinesterase and Butyrylcholinesterase - Important Enzymes of Human Body. *Acta Medica*, 47 (4), 215–228.
- Payumo, J. G., Jussaume, R., & Grimes, H. D. (2009). Protecting And Preserving Traditional Knowledge and Plant Genetic Resources : Is ASEAN There Yet? *Plant Genetic Resources : Characterization and Utilization*, 8 (1), 26–34.
- Pedrosa, T. do N., Barros, A. O., Nogueira, J. R., Fruet, A. C., Rodrigues, I. C., Calcagno, D. Q., Smith, M.A. C., Souza, T. P. D., Barros S. B. M.,

- Vasconcellos, M. C. D., da Silva, F. M. A. J., Koolen, H. H. F., Maria-Engler, S. & Lima, E. S. (2016). Anti-wrinkle and Anti-Whitening Effects of Juca (*Libidibia ferrea* Mart.) Extracts. *Archives of Dermatological Research*, 308(9), 643–654.
- Peñarrieta, J. M., Alvarado, J. A., Bergenståhl, B., & Åkesson, B. (2007). Spectrophotometric Methods for the Measurement of Total Phenolic Compounds and Total Flavonoids in Foods. *Revista Boliviana De Quimica*, 24 (1), 5–9.
- Pereira, D. M., Ferreres, F., Oliveira, J., Valentão, P., Andrade, P. B., & Sottomayor, M. (2009). Targeted Metabolite Analysis of *Catharanthus roseus* and its Biological Potential. *Food and Chemical Toxicology*, 47, 1349–1354.
- Pérez-Hernández, J., Zaldívar-Machorro, V. J., Villanueva-Porras, D., Vega-Ávila, E., & Chavarría, A. (2016). A Potential Alternative against Neurodegenerative Diseases. *Oxidative Medicine And Cellular Longevity*, 19 pages.
- Petharat, L., Wongsuphasawat, K., & Winther, K. (2015). Beneficial Effects of a Rose Hip Preparation Containing Seeds and Shells on Aging and Wrinkles of the Skin. *European Journal of Integrative Medicine*, 7, 46.
- Phipps, S. M., Sharaf, M. H. M., & Butterweck, V. (2007). Assessing Antioxidant Activity in Botanicals and other Dietary Supplements. *Pharmacopeial Forum*, 7, 6 pages.
- Pillai, N. S., Kannan, P. S., Vettivel, S. C., & Suresh, S. (2017). Optimization Of Transesterification of Biodiesel using Green Catalyst Derived from *Albizia Lebbeck* Pods by Mixture Design. *Renewable Energy*, 104, 185–196.
- Pimple, B. P., & Badole, S. L. (2013). Polyphenols: A Remedy for Skin Wrinkles. In R. R. Watson, V. R. Preedy, & S. Zibadi (Eds.). *Polyphenols in Human Health and Disease*. United Kingdom : Academic Press. pp. 861–869.
- Pinelo, M., Manzocco, L., Nunez, M. J., & Nicoli, M. C. (2004). Interaction among Phenols in Food Fortification: Negative Synergism on Antioxidant Capacity. *Journal of Agricultural and Food Chemistry*, 52, 1177–1180.
- Pires, T. C. S. P., Inês, M., Barros, L., & Ferreira, I. C. F. R. (2017). Nutritional and Chemical Characterization of Edible Petals and Corresponding Infusions : Valorization as New Food Ingredients. *Food Chemistry*, 220, 337–343.
- Pisano, M. B., Cosentino, S., Viale, S., Spanò, D., Corona, A., Esposito, F., Enzo Tramontano, E., Montoro, P., Tuberose, C. I. G., Medda, R. & Pintus, F. (2016).

Biological Activities of Aerial Parts Extracts of *Euphorbia characias*. *BioMed Research International*, 11pages.

Piowowski, J. P., Kiss, A. K., & Kozłowska-Wojciechowska, M. (2011). Anti-hyaluronidase and Anti-elastase Activity Screening of Tannin-Rich Plant Materials used in Traditional Polish Medicine for External Treatment of Diseases with Inflammatory Background. *Journal of Ethnopharmacology*, 137(1), 937–941.

Popa-wagner, A., Mitran, S., Sivanesan, S., Chang, E., & Buga, A. (2013). ROS and Brain Diseases: The Good, the Bad, and the Ugly. *Oxidative Medicine and Cellular Longevity*, 14 pages.

Porfírio, S., Falé, P. L. V., Madeira, P. J. A., Florêncio, M. H., Ascensão, L., & Serralheiro, M. L. M. (2010). Antiacetylcholinesterase and Antioxidant Activities of *Plectranthus barbatus* Tea, after *In Vitro* Gastrointestinal Metabolism. *Food Chemistry*, 122 (1), 179–187.

Prieto, M. A., Curran, T. P., Gowen, A., & Vázquez, J. A. (2015). An Efficient Methodology for Quantification Of Synergy and Antagonism in Single Electron Transfer Antioxidant Assays. *Food Research International*, 67, 284–298.

Prior, R. L., & Cao, G. (2000). Antioxidant Phytochemicals in Fruits and Vegetables: Diet and Health Implications. *HortScience*, 35, 588–592.

Queiros, B., Barreira, J. C. M., Sarmiento, A. C., & Ferreira, I. C. F. R. (2009). In Search of Synergistic Effects in Antioxidant Capacity of Combined Edible Mushrooms. *International Journal of Food Sciences and Nutrition*, 60 (S6), 160–172.

Rabe, J. H., Mamelak, A. J., Mcelgunn, P. J. S., Morison, W. L., & Sauder, D. N. (2006). Photoaging: Mechanisms and Repair. *Journal of the American Academy of Dermatology*, 55 (1), 1–19.

Rahal, A., Kumar, A., Singh, V., Yadav, B., Tiwari, R., Chakraborty, S., & Dhama, K. (2014). Oxidative stress, prooxidants, and antioxidants: The interplay. *BioMed Research International*, 19 pages.

Rao, P. V., Huey, L., Mohamed, S., Abdul-wahab, I. R., & Mei, S. (2016). Ethnomedicinal Knowledge of Temiar Ethnic Tribe of Lojing Highlands, Kelantan: A Source for Nutritional and Antioxidant Potential. *Proceedings of the 5th World Conference on Applied Sciences, Engineering and Technology*, 12–21.

- Re, R., Pellegrini, N., Proteggente, A., Pannala, A., Yang, M., & Rice-Evans, C. (1999). Antioxidant Activity Applying an Improved ABTS Radical Cation Decolorization Assay. *Free Radical Biology and Medicine*, 26(9), 1231–1237.
- Roriz, C. L., Barros, L., Carvalho, A. M., Santos-buelga, C., & Ferreira, I. C. F. R. (2015). Scientific Validation of Synergistic Antioxidant Effects in Commercialised Mixtures of *Cymbopogon citratus* and *Pterospartum tridentatum* or *Gomphrena globosa* for Infusions Preparation. *Food Chemistry*, 185, 16–24.
- Royer, M., Prado, M., García-Pérez, M. E., Diouf, P. N., & Stevanovic, T. (2013). Study of Nutraceutical, Nutricosmetics and Cosmeceutical Potentials of Polyphenolic Bark Extracts from Canadian Forest Species. *Pharma Nutrition*, 1(4), 158–167.
- Rubino, M. T., Maggi, D., Laghezza, A., Loiodice, F., Tortorella, P., & Moro, A. (2011). Identification of Novel Matrix Metalloproteinase Inhibitors by Screening Of Phenol Fragments Library. *Archive Der Pharmazie Chemistry in Life Science*, 344, 557–563.
- Sabran, S. F. (2016). *Documentation, Antimycobacterial Activity, and Phytochemical Profiling of Selected Medicinal Plants used by the Jakun Community in Johor*. Universiti Tun Hussein Onn Malaysia : Ph. D Thesis.
- Sabran, S. F., Mohamed, M., & Bakar, M. F. A. (2016). Ethnomedical Knowledge of Plants Used for the Treatment of Tuberculosis in Johor, Malaysia. *Evidence-Based Complementary and Alternative Medicine*, 2016, 12 pages.
- Sadeghia, Z., & Mahmood, A. (2014). Ethno-gynecological Knowledge of Medicinal Plants used by Baluch Tribes, Southeast of Baluchistan, Iran. *Brazilian Journal of Pharmacognosy*, 24 (6), 706–715.
- Sadowska-Bartosz, I., & Bartosz, G. (2014). Effects of Antioxidant Supplementation on Aging and Longevity. *BioMed Research International*, 17 pages.
- Salomone, S., Caraci, F., Leggio, G. M., Fedotova, J., & Drago, F. (2011). New Pharmacological Strategies for Treatment of Alzheimer's Disease : Focus on Disease Modifying Drugs. *British Journal of Clinical Pharmacology*, 73 (4), 504–517.
- Samuel, A. J. S. J., Kalusalingam, A., Chellappan, D. K., Gopinath, R., Radhamani, S., Husain, H. A., Muruganandham, V. & Promwichit, P. (2010). Ethnomedical Survey Of Plants used by the Orang Asli in Kampung Bawong, Perak, West

- Malaysia. *Journal of Ethnobiology and Ethnomedicine*, 6 (5), 6 pages.
- Sani, H. L., Malami, I., Hassan, S. W., Alhassan, A. M., Halilu, M. E. & Muhammad, A. (2015). Effects of Standardized Stem Bark Extract of *Mangifera Indica* L. in Wistar Rats with 2,4-Dinitrophenylhydrazine-Induced Haemolytic Anemia. *Pharmacognosy Journal*, 7(2), 89–96.
- Scheffe, H. (1963). The Simplex-Centroid Design for Experiments with Mixtures. *Journal of The Royal Statistical Society*, 25 (2), 235–263.
- Şenol, F. S., Orhan, I., Yilmaz, G., Çiçek, M., & Şener, B. (2010). Acetylcholinesterase, Butyrylcholinesterase, and Tyrosinase Inhibition Studies and Antioxidant Activities of 33 *Scutellaria* L. taxa from Turkey. *Food and Chemical Toxicology*, 48(3), 781–788.
- Shah, A. A., Dar, T. A., Dar, P. A., & Showkat A. Ganie, M. A. K. (2017). A Current Perspective on the Inhibition of Cholinesterase by Natural and Synthetic Inhibitors. *Current Drug Metabolism*, 18(2), 96–111.
- Shah, H., & Rawal, S. (2014). Screening of Topical Gel Containing Lycopene and Dexamethasone against UV Radiation Induced Photoaging In Mice. *Biomedicine and Aging Pathology*, 4(4), 303–308.
- Shakir, T., Coulibaly, A. Y., & Kehoe, P. G. (2013). An Exploration Of The Potential Mechanisms and Translational Potential of Five Medicinal Plants for Applications in Alzheimer ' S Disease. *American Journal of Neurodegenerative Disease*, 2(2), 70–88.
- Shin, N. H., Ryu, S. Y., Choi, E. J., Kang, S. H., Chang, I. M., Min, K. R., & Kim, Y. (1998). Oxyresveratrol as the Potent Inhibitor on DOPA Oxidase Activity of Mushroom Tyrosinase. *Biochemical and Biophysical Research Communications*, 243, 801–803.
- Si, H., & Liu, D. (2014). Dietary Antiaging Phytochemicals and Mechanisms Associated with Prolonged Survival. *Journal of Nutritional Biochemistry*, 25(6), 581–591.
- Sies, H. (1985). Oxidative Stress: Introductory Remarks. In Sies, H. *Oxidative Stress*. Orlando: Academic Press. pp. 1–8.
- Silva, T., Reis, J., Teixeira, J., & Borges, F. (2014). Alzheimer's Disease, Enzyme Targets and Drug Discovery Struggles: From Natural Products to Drug Prototypes. *Ageing Research Reviews*, 15, 116–145.
- Sindhuja, P. R., Prabhakaran, G., & Gokulshankar, S. (2014). Synergistic Effect of

- Anti-Oxidant, Anti-Tyrosinase and Anti-Bacterial Activities of *Tridax procumbens*, *Lantana camara*, *Euphorbia hirta* and *Thevetia peruviana* Plant Extracts for Cosmetic and Personal Care Applications. *International Journal of Pharmacy and Pharmaceutical Sciences*, 6(9), 7–10.
- Singhal, R. (2005). Medicinal Plants and Primary Health Care : The Role of Gender. *Journal of Health Management*, 7 (2), 277–293.
- Singleton, V. L., & Rossi, J. A. (1965). Colorimetry of Total Phenolics with Phosphomolybdic-Phosphotungstic Acid Reagents. *American Journal of Enology and Viticulture*, 16, 144–153.
- Smith, A. D., & Cuello, A. C. (1984). Alzheimer's Disease and Acetylcholinesterase-containing Neurons. *Lancet*, 1, 513.
- Smith, W. F. (2005). *Experimental Design For Formulation*. ASA-SIAM Series on Statistics and Applied Probability. Philadelphia : American Statistical Association.
- Soares, P. K., Bruns, R. E., & Scarminio, I. S. (2008). Statistical Mixture Design — Varimax Factor Optimization for Selective Compound Extraction from Plant Material. *Analytical Chimica Acta*, 3, 48–55.
- Solanki, I., Parihar, P., & Parihar, M. S. (2016). Neurodegenerative Diseases : From Available Treatments to Prospective Herbal Therapy. *Neurochemistry International*, 95, 100–108.
- Souza, P. M., Elias, S. T., Simeoni, L. A., Paula, J. E. de, Gomes, S. M., Guerra, E. N. S., Fonseca, Y. M., Silva, C., Silveira, D & Magalhaes, P. O. (2012). Plants from Brazilian Cerrado with Potent Tyrosinase Inhibitory Activity. *PLOS ONE*, 7 (11), 1–7.
- Spencer, J. P. E., Abd, M. M., Mohsen, E., Minihane, A., & Mathers, J. C. (2008). Biomarkers of The Intake of Dietary Polyphenols : Strengths, Limitations and Application in Nutrition Research. *British Journal of Nutrition*, 99, 12–22.
- Sung, J.-H., Park, S.-H., Seo, D.-H., Lee, J.-H., Hong, S.-W., & Hong, S.-S. (2009). Antioxidative and Skin-Whitening Effect of an Aqueous Extract of *Salicornia herbacea*. *Bioscience, Biotechnology, and Biochemistry*, 73(3), 552–556.
- Szwajgier, D. (2015). Anticholinesterase Activity of Selected Phenolic Acids and Flavonoids - Interaction Testing in Model Solutions. *Annals of Agricultural and Environmental Medicine*, 22(4), 690–694.

- Taeymans, J., Clarys, P., & Barel, A. O. (2014). Use of Food Supplements as Nutricosmetics in Health and Fitness. In Barel, A. O., Paye, M., Maibach & H. I. *Handbook of Cosmetic Science and Technology*. 4th ed. United State : CRC Press. pp. 583–596..
- Tamrakar, S., Tran, H. B., Nishida, M., Kaifuchi, S., Suhara, H., Doi, K., Fukami, K., Shimizu, G.P.P., Simizu, K. (2016). Antioxidative Activities of 62 Wild Mushrooms from Nepal and the Phenolic Profile of Some Selected Species. *Journal of Natural Medicines*, 11 pages.
- Tan, Y. P. & Chan, E. W. C. (2014). Antioxidant, Antityrosinase and Antibacterial Properties of Fresh and Process Leaves of *Anacardium occidentale* and *Piper betle*. *Food Bioscience*, 6, 17-23.
- Taofiq, O., Gonzalez-paramás, A. M., Martins, A., Filomena, M., & Ferreira, I. C. F. R. (2016). Mushrooms Extracts and Compounds in Cosmetics, Cosmeceuticals and Nutricosmetics - A Review. *Industrial Crop and Products*, 90, 38–48.
- Tauro, M., Laghezza, A., Loiodice, F., Piemontese, L., Capelli, D., Montanari, R., Pochetti, G., Pizio, A. D., Agamennone, M., Campestre, C. & Tortorella, P. (2016). Catechol-Based Matrix Metalloproteinase Inhibitors with Additional Antioxidative Activity. *Journal of Enzyme Inhibition and Medicinal Chemistry*, 31 (S4), 25–37.
- Taylor, C. E., & Wong, K. M. (1987). Some Aspects Of Herbal Medicine Among The Orang Hulu Community of Kampung Peta, Johore, Malaysia. *The Malayan Nature Journal*, 41, 317–328.
- Thaipong, K., Boonprakob, U., Crosby, K., Cisneros-Zevallos, L., & Hawkins Byrne, D. (2006). Comparison Of ABTS, DPPH, FRAP, and ORAC Assays for Estimating Antioxidant Activity from Guava Fruit Extracts. *Journal of Food Composition and Analysis*, 19(6), 669–675.
- Thring, T. S. A. (2012). *Method development and application for the assessment of the antiageing and antioxidant properties of herbal remedies*. Kingston University : Ph. D Thesis.
- Thring, T. S. A., Hili, P., & Naughton, D. P. (2009). Anti-Collagenase, Anti-Elastase and Anti-oxidant Activities of Extracts from 21 Plants. *BMC Complementary and Alternative Medicine*, 9 (27), 1–11.
- Tobin, D. J. (2017). Introduction to Skin Aging. *Journal of Tissue Viability*, 26 (1), 37–46.

- Tsao, R. (2015). Synergistic Interactions between Antioxidants used in Food Preservation. In Shahidi, F. *Handbook of Antioxidants for Food Preservation*. United Kingdom : Woodhead Publishing. pp. 335-347
- Tuler, A. C., & Nina, C. B. (2014). Women's Ethnomedicinal Knowledge in the Rural Community of São José da Figueira, Durandé, Minas Gerais, Brazil. *Revista Brasileira de Farmacognosia*, 24(2), 159–170.
- Van Overwalle, G. (2005). Protecting and Sharing Biodiversity and Traditional Knowledge: Holder and User Tools. *Ecological Economics*, 53(4), 585–607.
- Velioglu, Y. S., Mazza, G., Gao, L., & Oomah, B. D. (1998). Antioxidant Activity and Total Phenolics in Selected Fruits, Vegetables, and Grain Products. *Journal of Agricultural and Food Chemistry*, 46(10), 4113–4117.
- Viegas, C., Bolzani, V. D. S., Barreiro, E. J., & Fraga, C. A. M. (2005). New Anti-Alzheimer Drugs from Biodiversity: The Role of the Natural Acetylcholinesterase Inhibitors. *Mini Reviews in Medicinal Chemistry*, 5(10), 915–926.
- Villaflores, O. B., Chen, Y., Chen, C., & Yeh, J. (2012). Curcuminoids and Resveratrol as Anti-Alzheimer Agents. *Taiwanese Journal of Obstetrics & Gynecology*, 51(4), 515–525.
- Vinutha, B., Prashanth, D., Salma, K., Sreeja, S. L., Pratiti, D., Padmaja, R., Radhika, S., Amit, A., Venkateshwarlu, K., Deepak, M. (2007). Screening of Selected Indian Medicinal Plants for Acetylcholinesterase Inhibitory Activity. *Journal of Ethnopharmacology*, 109, 359–363.
- Wang, D., Solla, S. R. de, Lebeuf, M., Bisbicos, T., Barrett, G. C., & Alae, M. (2017). Determination of Linear and Cyclic Volatile Methylsiloxanes in Blood of Turtles, Cormorants, and Seals From Canada. *Science of the Total Environment*, 574, 1254–1260.
- Wang, F., Zhao, S., Li, F., Zhang, B., Qu, Y., Sun, T., Luo, T., Li, D. (2014). Investigation of Antioxidant Interactions between *Radix Astragali* and *Cimicifuga foetida* and Identification Of Synergistic Antioxidant Compounds. *PLOS ONE*, 9 (1), 1-12.
- Wang, S., Meckling, K. A., Marcone, M. F., Kakuda, Y., & Tsao, R. (2011). Synergistic, Additive, and Antagonistic Effects of Food Mixtures on Total Antioxidant Capacities. *Journal of Agricultural and Food Chemistry*, 59(3), 960–968.

- Weinert, B. T., & Timiras, P. S. (2003). Physiology of Aging. Invited review: Theories of Aging. *Journal of Applied Physiology*, 95, 1706–1716.
- Weldegerima, B. (2009). Review on the Importance of Documenting Ethnopharmacological Information on Medicinal Plants. *African Journal of Pharmacy and Pharmacology*, 3(9), 400–403.
- WHO. (2000). *General guidelines for methodologies on research and evaluation of traditional medicine*. Geneva: World Health Organization.
- WHO. (2003). Traditional Medicine. Retrieved April 19, 2017, from <http://www.who.int/mediacentre/factsheets/2003/fs134/en/>
- Wightman, E. L. (2017). Potential Benefits of Phytochemicals against Alzheimer's Disease. *Proceedings of the Nutrition Society Proceedings*, 76 (2), 106-112.
- WIPO. (2001). *Intellectual property needs and expectations of traditional knowledge holders. WIPO Report on Fact-finding Missions on Intellectual Property and Traditional Knowledge (1998-1999)*.
- WIPO. (2012). *The World Intellectual Property Organization Traditional Knowledge Documentation Toolkit. Consultation draft*.
- Wu, L., Jou, A. F., Chen, S., Tien, C., Cheng, C., Fan, N., & Ho, J. A. (2010). Antioxidant, Anti-inflammatory And Anti-browning Activities of Hot Water Extracts of Oriental Herbal Teas. *Food and Function*, 1, 200–208.
- Xu, X., Li, F., Zhang, X., Li, P., Zhang, X., Wu, Z., & Li, D. (2014). *In Vitro* Synergistic Antioxidant Activity and Identification of Antioxidant Components from *Astragalus membranaceus* and *Paeonia lactiflora*. *PLOS ONE*, 9 (5), 1-9.
- Yang, J., & Liu, R. H. (2013). The Phenolic Profiles and Antioxidant Activity in Different Types Of Tea. *International Journal of Food Science and Technology*, 48 (1), 163–171.
- Yang, W.-J., Li, D.-P., Li, J.-K., Li, M.-H., Chen, Y.-L., & Zhang, P.-Z. (2009). Synergistic Antioxidant Activities of Eight Traditional Chinese Herb Pairs. *Biological and Pharmaceutical Bulletin*, 32(6), 1021–1026.
- Yang, Z., Zhang, D., Ren, J., Yang, M., & Li, S. (2012). Acetylcholinesterase Inhibitory Activity of the Total Alkaloid from Traditional Chinese Herbal Medicine for Treating Alzheimer's disease. *Medicinal Chemistry Research*, 21(6), 734–738.
- Yazbek, P. B., Tezoto, J., Cassas, F., & Rodrigues, E. (2015). Plants used during Maternity, Menstrual Cycle and Other Women's Health Conditions among

- Brazilian Cultures. *Journal of Ethnopharmacology*, 179, 310–31.
- Yoshiara, L. Y., Madeira, T. B., Delaroza, F., Silva, C. I. O. D. A., Ida, E. I., & Bonifa, J. (2012). Optimization of Soy Isoflavone Extraction With Different Solvents using the Simplex-Centroid Mixture Design. *International Journal of Food Science and Nutrition*, 63(8), 978–986.
- Yu, X., Zhao, M., Liu, F., Zeng, S., & Hu, J. (2013). Identification of 2,3-dihydro-3,5-dihydroxy-6-methyl-4H-pyran-4-one as a Strong Antioxidant in Glucose – Histidine Maillard Reaction Products. *Food Research International*, 51, 397–403.
- Zengin, G., Nithiyanantham, S., Locatelli, M., Ceylan, R., Uysal, S., Aktumsek, A., Selvi, P. K. & Maskovic, P. (2016). Screening of *In Vitro* Antioxidant and Enzyme Inhibitory Activities of Different Extracts from Two Uninvestigated Wild Plants: *Centranthus longiflorus* subsp. *longiflorus* and *Cerithe minor* subsp. *auriculata*. *European Journal of Integrative Medicine*, 8(3), 286–292.
- Zheng, X., Wei, J., Sun, W., Li, R., Liu, S., & Dai, H. (2013). Ethnobotanical Study On Medicinal Plants around Limu Mountains of Hainan Island, China. *Journal of Ethnopharmacology*, 148(3), 964–974.
- Zhishen, J., Tang, M., & Wu, J. (1999). The Determination of Flavonoid Contents in Mulberry and their Scavenging Effects on Superoxide Radicals. *Food Chemistry*, 64, 555–559.
- Zielinski, A. A. F., Granato, D., Alberti, A., Nogueira, A., Demiate, I. M., & Haminiuk, C. W. I. (2015). Modelling the Extraction of Phenolic Compounds and *In Vitro* Antioxidant Activity of Mixtures of Green, White and Black Teas (*Camellia sinensis* L. Kuntze). *Journal of Food Science and Technology*, 52 (11), 6966–6977.